

Partonic Pole Matrix Elements for Fragmentation

S. Meißner

Institut für Theoretische Physik II, Ruhr-Universität Bochum, 44780 Bochum, Germany

A. Metz

Department of Physics, Temple University, Philadelphia, Pennsylvania 19122-6082, USA

(Dated: 5 May 2009)

A model-independent analysis of collinear three-parton correlation functions for fragmentation is performed. By investigating their support properties it is shown, in particular, that the so-called partonic pole matrix elements vanish. This sheds new light on the understanding of transverse single spin asymmetries in various hard semi-inclusive reactions. Moreover, it gives additional strong evidence for the universality of transverse-momentum-dependent fragmentation functions.

PACS numbers: 12.38.Bx, 12.39.St, 13.85.Ni, 13.87.Fh

I. Introduction and definitions.—QCD factorization theorems separate the cross section for a large number of hard processes into a perturbatively calculable part as well as nonperturbative correlation functions [1]. These correlation functions are parton distributions (PDFs), fragmentation functions (FFs), etc. [2, 3]. So far the main focus was on the leading (twist-2) two-parton correlators, and in the meantime an enormous amount of information about these objects has been obtained.

However, there is a set of interesting phenomena which—in the typically used collinear picture with partons moving in the direction of their parent hadrons—cannot be described by means of twist-2 correlators. Perhaps the most intriguing observables among those are transverse single spin asymmetries (SSAs) for which effects of up to 40 % were observed in hard hadronic reactions such as $p^\uparrow p \rightarrow \pi X$ and $\bar{p}^\uparrow p \rightarrow \pi X$ at FermiLab [4, 5] and more recently also at the Relativistic Heavy Ion Collider [6, 7, 8, 9]. In a factorized QCD approach one can generate such asymmetries by using particular collinear twist-3 three-parton correlators. This idea was brought up in [10] and worked out in more detail in [11] (see also Refs. [12, 13]). If one of the three partons becomes soft, one hits the pole of a particle propagator in the hard part of a process providing an imaginary part (a nontrivial phase) which, quite generally, can result in SSAs [10, 11]. Therefore, multiparton correlators with one (or more) soft partons are called gluonic pole or fermionic pole matrix elements, depending on whether the soft parton is a gluon or a quark.

Already a lot of work on different aspects of these partonic pole matrix elements (PPMEs) exists, which has considerably improved our understanding about such objects and the physics underlying SSAs. So far the papers in the field of PPMEs were mostly dealing with correlators that describe the parton structure of the target (PPMEs for parton distributions). The knowledge about multiparton correlators for fragmentation, however, is still rather poor, and it is the aim of the present Letter to give new insights into these objects.

Nonzero PPMEs for fragmentation would contribute to a large number of transverse SSAs. By using simple models for such three-parton matrix elements pioneering phenomenological studies on the processes $p^\uparrow p \rightarrow \pi X$, $pp \rightarrow \Lambda^\uparrow X$, $ep^\uparrow \rightarrow e\pi X$, and $ep \rightarrow e\Lambda^\uparrow X$ were carried out in [14, 15]. Other work discussed PPMEs for fragmentation in connection with $p^\uparrow p \rightarrow \pi\pi X$, $p^\uparrow p \rightarrow \text{jet } \pi X$ [16, 17], as well as $pp \rightarrow (\Lambda^\uparrow \text{jet}) \text{jet } X$ with a Λ hyperon detected in a jet [18]. Note that this already long list of semi-inclusive reactions, which might be related to PPMEs for fragmentation, is not yet complete.

Despite the potential importance of PPMEs for fragmentation it was, however, so far not even clear whether they exist. In fact, recent spectator model calculations imply (show) that these objects vanish [19, 20, 21]. In this Letter it is proven for the first time in a fully model-independent way that PPMEs for fragmentation vanish, which constitutes the main outcome of our work. To obtain this result we analyze the support properties of collinear twist-3 three-parton fragmentation correlators.

Now in order to be more specific we consider essentially two kinds of correlators in fragmentation: the quark-gluon-quark (qgq) correlator as well as the gluon-gluon-gluon (ggg) correlator. The qgq correlator (see also Fig. 1) is defined by

$$\begin{aligned} \Delta_F^i(\tfrac{1}{z}, \tfrac{1}{z'}) &= \sum_X \int \frac{d\xi^+}{2\pi} \frac{d\eta^+}{2\pi} e^{i\frac{1}{z'}P^-\xi^+ + i(\frac{1}{z} - \frac{1}{z'})P^-\eta^+} \\ &\times \frac{1}{3} \text{Tr} \left[\langle 0 | g t_a F_a^{-i}(\eta^+) \psi(\xi^+) | P, X \rangle \right. \\ &\left. \times \langle P, X | \bar{\psi}(0^+) | 0 \rangle \right], \end{aligned} \quad (1)$$

where $v^\pm = (v^0 \pm v^3)/\sqrt{2}$ and $\vec{v}_T = (v^1, v^2)$ are the light-cone components of a generic 4-vector $v = (v^0, v^1, v^2, v^3)$ and P^- in Eq. (1) is the (large) minus momentum of the hadron in the final state. In Fig. 1 the minus momenta of the three partons are given. The correlator in Eq. (1) is a matrix in Dirac space, and the trace is acting in color space. We make use of the light-cone gauge $A^- = 0$, in which Wilson lines between the field operators reduce to




FIG. 1: Kinematics of the qgq correlator in Eq. (1). The wiggly line with the open circle indicates the component F_a^{-i} of the gluon field strength tensor.

the unit matrix. Note that replacing the quark fields ψ in Eq. (1) by the corresponding charge conjugated quark fields ψ^C defines the $\bar{q}g\bar{q}$ correlator $\hat{\Delta}_P^i$. The ggg correlator containing three field strength tensors (see also Fig. 2) is given by

$$\begin{aligned} \hat{\Gamma}_{F,f}^{i,jk}(\frac{1}{z}, \frac{1}{z'}) &= \sum_X \int \frac{d\xi^+}{2\pi} \frac{d\eta^+}{2\pi} e^{i\frac{1}{z'}P^-\xi^+ + i(\frac{1}{z} - \frac{1}{z'})P^-\eta^+} \\ &\times i f_{abc} \langle 0 | g F_b^{-j}(\eta^+) F_c^{-k}(\xi^+) | P, X \rangle \\ &\times \langle P, X | F_a^{-i}(0^+) | 0 \rangle, \end{aligned} \quad (2)$$

with f_{abc} denoting the (antisymmetric) structure constants of the $SU(3)$ group. If in Eq. (2) the symmetric structure constants d_{abc} are used instead of the $i f_{abc}$ one obtains the additional, independent ggg correlator $\hat{\Gamma}_{F,d}^{i,jk}$.

We point out that the objects defined above form a complete set of collinear twist-3 three-parton correlators for fragmentation. Two other sets can be defined by either substituting the gluon field strength tensor F^{-i} at position η^- with the covariant derivative D_T^i or by shifting it from the first to the second matrix element in Eqs. (1) and (2). These sets are, however, not independent of the one we are using. In the first case they are connected in analogy to Ref. [15] while in the second case they are just the Hermitian conjugates of each other.

In the following we will show that the correlators in Eqs. (1) and (2) vanish if the longitudinal (minus) momentum of one of the two partons on the right-hand




FIG. 2: Kinematics of the ggg correlator in Eq. (2). The wiggly lines with the open circles indicate the different components of the gluon field strength tensor.

side (rhs) of the cut in Figs. 1 and 2 vanishes. Precisely these kinematical conditions are what defines the PPMEs.

II. Support properties of three-parton correlators.— We outline our analysis for the case of the qgq correlator in Eq. (1). For this correlator we obtain by inserting a complete set of intermediate states $|Y\rangle$ and by using the translation properties of field operators

$$\begin{aligned} \Delta_F^i(\frac{1}{z}, \frac{1}{z'}) &= \sum_{X,Y} \int \frac{d\xi^+}{2\pi} \frac{d\eta^+}{2\pi} e^{i\frac{1}{z'}P^-\xi^+ + i(\frac{1}{z} - \frac{1}{z'})P^-\eta^+} \\ &\times \frac{1}{3} \text{Tr} \left[\langle 0 | g t_a F_a^{-i}(\eta^+) | Y \rangle \right. \\ &\times \langle Y | \psi(\xi^+) | P, X \rangle \langle P, X | \bar{\psi}(0^+) | 0 \rangle \Big] \\ &= \sum_{X,Y} \delta \left((\frac{1}{z} - 1)P^- - \sum_i p_i^- \right) \\ &\times \delta \left((\frac{1}{z} - \frac{1}{z'})P^- - \sum_j q_j^- \right) \\ &\times \frac{1}{3} \text{Tr} \left[\langle 0 | g t_a F_a^{-i}(0^+) | Y \rangle \right. \\ &\times \langle Y | \psi(0^+) | P, X \rangle \langle P, X | \bar{\psi}(0^+) | 0 \rangle \Big]. \end{aligned} \quad (3)$$

Here p_i and q_j denote the momenta of the particles in the intermediate states $|X\rangle$ and $|Y\rangle$, respectively. As all particles in the intermediate states are on-shell, their momenta have to satisfy $p_i^2 = m_i^2$ and $p_i^0 \geq 0$ as well as $q_j^2 = m_j^2$ and $q_j^0 \geq 0$, which leads to $p_i^- \geq 0$ and $q_j^- \geq 0$. This means, however, that the δ functions in Eq. (3) only give a nonzero contribution if

$$\frac{1}{z} \geq 1 \quad \text{and} \quad \frac{1}{z} \geq \frac{1}{z'}. \quad (4)$$

The case $\frac{1}{z} = \frac{1}{z'}$ is now of particular interest because it corresponds to a PPME. From Eq. (3) one finds that here the second δ function only contributes if $q_j^- = 0$ for all j , which leads to $q_j^2 = -\bar{q}_{jT}^2 \leq 0$. Combined with the on-shell condition $q_j^2 = m_j^2 \geq 0$ it immediately follows that all particles in the intermediate state $|Y\rangle$ have to be massless and move only in the plus direction, i.e., $q_j^- = \bar{q}_{jT} = 0$ for all j , in order to obtain a nonvanishing contribution to the qgq correlator. In this case, however, the matrix element

$$M^{-i}(q_j) = \langle 0 | g t_a F_a^{-i}(0^+) | Y \rangle \quad (5)$$

vanishes. This matrix element is an antisymmetric Lorentz tensor, which can be expressed in terms of the Lorentz vectors it depends on—the momenta q_j and the polarization vectors $\epsilon(q_j)$ of the massless particles in the intermediate state—multiplied by some scalar functions. One possible decomposition of the matrix element in Eq. (5) with general Lorentz indices is

$$\begin{aligned} M^{\mu\nu}(q_j) &= \sum_{m,n} \left[q_m^\mu q_n^\nu A_{mn}(q_j) + q_m^\mu \epsilon^\nu(q_n) B_{mn}(q_j) \right. \\ &\quad \left. + \epsilon^\mu(q_m) \epsilon^\nu(q_n) C_{mn}(q_j) - \{\mu \leftrightarrow \nu\} \right]. \end{aligned} \quad (6)$$

However, for massless, on-shell particles moving in the plus direction the minus components of the polarization vectors $\epsilon(q_j)$ have to vanish because they correspond to some combination of the unphysical timelike and longitudinal polarization components. Therefore, none of the terms in Eq. (6) can contribute to the matrix element M^{-i} in Eq. (5) as we have $q_j^- = \vec{q}_{jT}^- = \epsilon^-(q_j) = 0$ for all j . This means that finally the qqq correlator in Eq. (3) vanishes for $\frac{1}{z} = \frac{1}{z'}$ and we only get a nonzero contribution if

$$\frac{1}{z} \geq 1 \quad \text{and} \quad \frac{1}{z} > \frac{1}{z'}. \quad (7)$$

On the other hand, if we permute the quark and the gluon field in the first matrix element in Eq. (1) before we insert the complete set of intermediate states $|Y\rangle$ into the qqq correlator, we find

$$\begin{aligned} \Delta_F^i(\frac{1}{z}, \frac{1}{z'}) &= \sum_{X,Y} \delta\left((\frac{1}{z}-1)P^- - \sum_i p_i^-\right) \\ &\times \delta\left(\frac{1}{z'}P^- - \sum_j q_j^-\right) \\ &\times \frac{1}{3} \text{Tr} \left[\langle 0 | t_a \psi(0^+) | Y \rangle \right. \\ &\left. \times \langle Y | g F_a^{-i}(0^+) | P, X \rangle \langle P, X | \bar{\psi}(0^+) | 0 \rangle \right]. \end{aligned} \quad (8)$$

In this case an analysis of the δ functions analogous to the one for the δ functions in Eq. (3) yields, that the qqq correlator only differs from zero if

$$\frac{1}{z} \geq 1 \quad \text{and} \quad \frac{1}{z'} > 0. \quad (9)$$

Here $\frac{1}{z'} = 0$, which also corresponds to a PPME, can be excluded because the matrix element $\langle 0 | t_a \psi(0^+) | Y \rangle$ vanishes if all particles in the intermediate state $|Y\rangle$ are massless. Therefore, from Eqs. (7) and (9) we conclude that the only nonzero contribution to the qqq correlator in Eq. (1) comes from

$$0 \leq z \leq 1 \quad \text{and} \quad 0 < \frac{z}{z'} < 1. \quad (10)$$

The same results also hold for the $\bar{q}g\bar{q}$ correlator and the two ggg correlators. For the PPMEs this leads to the model-independent results

$$\Delta_F^i(\frac{1}{z}, \frac{1}{z}) = \Delta_F^i(\frac{1}{z}, 0) = 0, \quad (11)$$

$$\bar{\Delta}_F^i(\frac{1}{z}, \frac{1}{z}) = \bar{\Delta}_F^i(\frac{1}{z}, 0) = 0, \quad (12)$$

$$\hat{\Gamma}_{F,f/d}^{i,jk}(\frac{1}{z}, \frac{1}{z}) = \hat{\Gamma}_{F,f/d}^{i,jk}(\frac{1}{z}, 0) = 0. \quad (13)$$

We repeat that the PPMEs for fragmentation, therefore, cannot be responsible for single spin asymmetries as they all vanish according to Eqs. (11)–(13).

Note that the support properties of three-parton correlators on the distribution side can be found by a corresponding analysis. In that case, however, PPMEs do not vanish which agrees with the well-established existence of

these objects. The key difference between both analyses is the absence of the vacuum state on the PDF side.

III. Universality of transverse-momentum-dependent fragmentation functions. — The vanishing of gluonic pole matrix elements (GPMEs) for fragmentation is intimately connected with the universality of transverse-momentum-dependent two-parton fragmentation functions (TMD FFs) [17, 22]. To illustrate the relation in more detail we consider the correlator

$$\begin{aligned} \Delta^{[U]}(\frac{1}{z}, \vec{k}_T) &= \sum_X \int \frac{d\xi^+}{2\pi} \frac{d^2\vec{\xi}_T}{(2\pi)^2} e^{ik \cdot \xi} \frac{1}{3} \text{Tr} \left[\langle 0 | \mathcal{W}^{[U]}(0, \xi) \right. \\ &\left. \times \psi(\xi) | P, X \rangle \langle P, X | \bar{\psi}(0) | 0 \rangle \right]_{\xi^- = 0}, \end{aligned} \quad (14)$$

which defines TMD FFs when appropriate traces in Dirac space are taken [23, 24]. At leading twist there exist eight FFs. (We restrict the explicit discussion here to the quark FFs, but our general conclusions apply to gluon FFs as well.) The TMD FFs depend both on the longitudinal momentum $k^- = P^-/z$ and the transverse momentum $\vec{k}_T$ (relative to the detected hadron) of the fragmenting quark. The Wilson line $\mathcal{W}^{[U]}$ cannot be neglected in any gauge [25, 26], and its path $\mathcal{U}(0; \xi)$ is determined by the physical process under consideration [22, 27, 28]. More details about the precise definition of TMD correlators and, in particular, the choice of Wilson lines can be found in [29, 30, 31] and references therein.

The path dependence of the correlator in Eq. (14) implies a potential nonuniversality of TMD FFs. For instance, in semi-inclusive deep inelastic scattering (DIS) one has, *a priori*, past-pointing Wilson lines, while they are future pointing in e^+e^- annihilation. In Ref. [20] it was argued, however, that factorization can be established such that TMD FFs have the same Wilson lines in both processes. But the analysis in [20] made use of a spectator model and was not carried out to arbitrary order in perturbation theory. As a consequence, in the community doubts remained concerning the generality of this result.

In the following we consider k_T moments of the correlator in Eq. (14). We define the zeroth moment by

$$\Delta(\frac{1}{z}) = \int d^2\vec{k}_T \Delta^{[U]}(\frac{1}{z}, \vec{k}_T). \quad (15)$$

In this case the k_T integration eliminates all the path dependence implying, in particular, that the three leading twist collinear FFs [23, 24], which are given by Δ in Eq. (15), are universal. For the more interesting case of the first k_T moment of the correlator in Eq. (14) one finds [17, 22]

$$\begin{aligned} \Delta_{\partial}^{i[U]}(\frac{1}{z}) &= \int d^2\vec{k}_T k_T^i \Delta^{[U]}(\frac{1}{z}, \vec{k}_T) \\ &= \tilde{\Delta}_{\partial}^i(\frac{1}{z}) + C_F^{[U]} \pi \Delta_F^i(\frac{1}{z}, \frac{1}{z}). \end{aligned} \quad (16)$$

Here $\tilde{\Delta}_g^i$ is the universal part, while the second term on the rhs in Eq. (16) contains all the potential nonuniversal behavior of the correlator. The latter is given by a GPME multiplied with the calculable and path-dependent so-called gluonic pole factor $C_F^{[M]}$. According to Eq. (11) the GPME vanishes, and, therefore, we conclude that the first moment of the TMD correlator in Eq. (14) is universal as well. This leads to universal second k_T moments of four specific leading twist TMD FFs, where currently the Collins FF represents the most important one. This universality is a prerequisite for a combined analysis of transverse-momentum-dependent data from semi-inclusive DIS and e^+e^- annihilation [32, 33], and, in particular, for the first extraction of the transversity distribution of the nucleon [33].

We repeat that universality of TMD FFs was already obtained previously in low order spectator model calculations [19, 20, 34]. However, so far no model-independent proof has been given. Therefore, our result, even though the present analysis is restricted to certain k_T moments of FFs, provides the first strong model-independent support for the universality of TMD FFs.

IV. Conclusions.—We have proved in a model-independent way that the so-called partonic pole matrix elements for fragmentation vanish. Therefore, in contrast to conjectures in the literature, transverse single spin asymmetries in hard semi-inclusive reactions are not related to such objects. Our finding does not mean that single spin asymmetries, in general, cannot be connected to collinear fragmentation correlators. This topic requires further investigation.

The vanishing of the partonic pole matrix elements also implies that certain k_T moments of transverse-momentum-dependent fragmentation functions are universal. This result confirms earlier studies, and represents the first fully general and model-independent proof of this kind. Hence, we now have additional strong evidence for the universality of transverse-momentum-dependent fragmentation functions. For the future one might hope that higher k_T moments of fragmentation functions can be analyzed along the lines presented in this Letter.

This work has been partially supported by the BMBF (Verbundforschung), the DFG, and the EU Integrated Infrastructure Initiative Hadronphysics Project under Contract No. RII3-CT-2004-506078.

-
- [1] J. C. Collins, D. E. Soper, and G. Sterman, *Adv. Ser. Direct. High Energy Phys.* **5**, 1 (1988).
 [2] J. C. Collins and D. E. Soper, *Nucl. Phys. B* **194**, 445 (1982).

- [3] R. L. Jaffe and X. Ji, *Nucl. Phys. B* **375**, 527 (1992).
 [4] D. L. Adams *et al.* (E704 Collaboration), *Phys. Lett. B* **261**, 201 (1991); *Phys. Lett. B* **264**, 462 (1991); *Z. Phys. C* **56**, 181 (1992).
 [5] A. Bravar *et al.* (E704 Collaboration), *Phys. Rev. Lett.* **77**, 2626 (1996).
 [6] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **92**, 171801 (2004).
 [7] S. S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **95**, 202001 (2005).
 [8] I. Arsene *et al.* (BRAHMS Collaboration), *Phys. Rev. Lett.* **101**, 042001 (2008).
 [9] B. I. Abelev *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **101**, 222001 (2008).
 [10] A. V. Efremov and O. V. Teryaev, *Sov. J. Nucl. Phys.* **36**, 140 (1982) [*Yad. Fiz.* **36**, 242 (1982)]; *Phys. Lett. B* **150**, 383 (1985).
 [11] J.-W. Qiu and G. Sterman, *Phys. Rev. Lett.* **67**, 2264 (1991); *Nucl. Phys. B* **378**, 52 (1992); *Phys. Rev. D* **59**, 014004 (1998).
 [12] C. Kouvavis, J.-W. Qiu, W. Vogelsang, and F. Yuan, *Phys. Rev. D* **74**, 114013 (2006).
 [13] H. Eguchi, Y. Koike, and K. Tanaka, *Nucl. Phys. B* **763**, 198 (2007).
 [14] Y. Koike, arXiv:hep-ph/0106260; *AIP Conf. Proc.* **675**, 449 (2003); *AIP Conf. Proc.* **675**, 574 (2003).
 [15] H. Eguchi, Y. Koike, and K. Tanaka, *Nucl. Phys. B* **752**, 1 (2006).
 [16] A. Bacchetta, C. J. Bomhof, P. J. Mulders, and F. Pijlman, *Phys. Rev. D* **72**, 034030 (2005).
 [17] C. J. Bomhof and P. J. Mulders, *JHEP* **0702**, 029 (2007).
 [18] D. Boer, C. J. Bomhof, D. S. Hwang, and P. J. Mulders, *Phys. Lett. B* **659**, 127 (2008).
 [19] A. Metz, *Phys. Lett. B* **549**, 139 (2002).
 [20] J. C. Collins and A. Metz, *Phys. Rev. Lett.* **93**, 252001 (2004).
 [21] L. P. Gamberg, A. Mukherjee, and P. J. Mulders, *Phys. Rev. D* **77**, 114026 (2008).
 [22] D. Boer, P. J. Mulders, and F. Pijlman, *Nucl. Phys. B* **667**, 201 (2003).
 [23] P. J. Mulders and R. D. Tangerman, *Nucl. Phys. B* **461**, 197 (1996) [Erratum-ibid. *B* **484**, 538 (1997)].
 [24] A. Bacchetta *et al.*, *JHEP* **0702**, 093 (2007).
 [25] X. Ji and F. Yuan, *Phys. Lett. B* **543**, 66 (2002).
 [26] A. V. Belitsky, X. Ji, and F. Yuan, *Nucl. Phys. B* **656**, 165 (2003).
 [27] J. C. Collins, *Phys. Lett. B* **536**, 43 (2002).
 [28] C. J. Bomhof, P. J. Mulders, and F. Pijlman, *Phys. Lett. B* **596**, 277 (2004).
 [29] J. C. Collins, arXiv:0808.2665 [hep-ph].
 [30] F. Hautmann and H. Jung, *AIP Conf. Proc.* **1056**, 79 (2008).
 [31] I. O. Cherednikov and N. G. Stefanis, arXiv:0811.4357 [hep-ph].
 [32] A. V. Efremov, K. Goeke, and P. Schweitzer, *Phys. Rev. D* **73**, 094025 (2006).
 [33] M. Anselmino *et al.*, *Phys. Rev. D* **75**, 054032 (2007).
 [34] F. Yuan, *Phys. Rev. Lett.* **100**, 032003 (2008); *Phys. Rev. D* **77**, 074019 (2008).